

Research Article

Efficacy of Alveolar Ridge Preservation after Maxillary Molar Extraction in Reducing Crestal Bone Resorption and Sinus Pneumatization: A Multicenter Prospective Case-Control Study

Teresa Lombardi,¹ Fabio Bernardello,² Federico Berton ,³
Davide Porrelli,³ Antonio Rapani,³ Alvise Camurri Piloni,³ Luca Fiorillo,⁴
Roberto Di Lenarda ,³ and Claudio Stacchi ³

¹Private Practice, Cassano allo Ionio (CS), Italy

²Private Practice, Terranegra di Legnago (VR), Italy

³Department of Medical, Surgical and Health Sciences, University of Trieste, Italy

⁴Department of Biomedical and Dental Sciences, Morphological and Functional Images, School of Dentistry University of Messina, Messina, Italy

Correspondence should be addressed to Claudio Stacchi; claudio@stacchi.it

Received 12 June 2018; Accepted 23 October 2018; Published 4 November 2018

Guest Editor: Tolga Tozum

Copyright © 2018 Teresa Lombardi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aim. To evaluate, with three-dimensional analysis, the effectiveness of alveolar ridge preservation (ARP) after maxillary molar extraction in reducing alveolar bone resorption and maxillary sinus pneumatization when compared to unassisted socket healing. **Methods.** Patients were included in the study following inclusion criteria and underwent minimally traumatic maxillary molar extraction followed by ARP using synthetic nanohydroxyapatite (Fisiograft Bone, Ghimas, Italy) (test group) or unassisted socket healing (control group). Cone-beam computerized tomographies (CBCT) were performed immediately after tooth extraction (T0) and 6 months postoperatively (T1). CBCTs were superimposed by using a specific software (Amira, Thermo Fisher Scientific, USA) and the following items were analyzed in both groups: (i) postextractive maxillary sinus floor expansion in coronal direction and (ii) postextractive alveolar bone dimensional changes (both vertical and horizontal). All data were tested for normality and equality of variance and subsequently analyzed by independent samples T-test and Mann–Whitney test. **Results.** Thirty patients were treated by three centers and twenty-six (test n=13; control n=13) were included in the final analysis. Mean sinus pneumatization at T1 was 0.69 ± 0.48 mm in the test group and 1.04 ± 0.67 mm in the control group ($p=0.15$). Mean vertical reduction of the alveolar bone at T1 was 1.62 ± 0.49 mm in the test group and 2.01 ± 0.84 mm in the control group ($p=0.08$). Mean horizontal resorption of crestal bone at T1 was 2.73 ± 1.68 mm in test group and 3.63 ± 2.24 mm in control group ($p=0.24$). **Conclusions.** It could be suggested that ARP performed after maxillary molar extraction may reduce the entity of sinus pneumatization and alveolar bone resorption, compared to unassisted socket healing. This technique could decrease the necessity of advanced regenerative procedures prior to dental implant placement in posterior maxilla.

1. Introduction

After dental extraction, the alveolar bone undergoes a remodeling process resulting in horizontal and vertical reduction of crestal dimensions [1, 2]. Ridge resorption may lead to inadequate bone volume for dental implants insertion and create both functional and esthetic issues during prosthetic rehabilitation [3]: in the posterior maxilla, in particular,

a substantial percentage of edentulous patients may need bone augmentation procedures to allow a proper implant placement and reach satisfactory results [4]. Many surgical solutions are currently available to regenerate an adequate amount of bone in the atrophic crests, including lateral and transcrestal sinus floor elevation, guided bone regeneration, and block grafting [5–10]. However, all of these options are associated with significant rate of complications, increased

morbidity, high costs, and prolonged time of therapy [11–13]. In the attempt to reduce the need for advanced surgical procedures and to simplify the treatment plan, specific surgical techniques were developed to reduce postextractive ridge resorption [14, 15]. Alveolar ridge preservation (ARP), with the application of different biomaterials, is the most common procedure aiming to control crestal bone resorption following dental extractions [16–18]. A recent systematic review confirmed that ARP results in significant reduction of vertical bone loss following dental extraction when compared to spontaneous socket healing, while its protective effect on horizontal reduction of the alveolar bone was found to be variable. Furthermore, no specific type of ARP could be demonstrated to be more effective than others in preventing alveolar crest shrinkage [19]. In a recent study, ARP performed in the posterior maxilla with a combination of allograft and collagen membrane resulted in 1.0 mm crestal height reduction and in approximately 2.5 mm loss of ridge width [20]; this volumetric contraction was lower than the one observed in extraction sites of the same area after spontaneous healing [21, 22].

Moreover, following tooth extraction in the posterior maxilla, postextractive crestal bone resorption may be associated with maxillary sinus pneumatization, which may contribute to a further decrease of the available bone volume for implant placement. The reasons for sinus pneumatization after tooth extraction are still debated and poorly understood: a possible explanation is a shift of the physiologic bone remodeling process towards a resorptive pattern, due to lack of functional forces which are normally transferred to the bone when the tooth is present [23]. This particular type of disuse atrophy occurs according to Wolff's law and is enhanced by the presence of a positive air pressure into the sinus cavity [24]. Previous human studies demonstrated a downward expansion of the maxillary sinus after dental extraction and showed that the expansion was larger if the extracted tooth was surrounded by a superiorly curved sinus floor [25, 26].

The effectiveness of ARP procedures in preventing maxillary sinus expansion has been recently evaluated in a retrospective study using bidimensional radiographs, showing significant differences in terms of postextractive pneumatization between test (ARP using bovine derived xenograft) and control group (spontaneous healing) [26].

The aim of this multicenter prospective case-control study was to evaluate, with three-dimensional analysis, the clinical effectiveness of ARP after maxillary molar extraction in reducing alveolar ridge resorption and maxillary sinus pneumatization when compared to unassisted socket healing.

2. Materials and Methods

2.1. Study Design. This multicenter prospective case-control study has been designed and conducted in accordance with the Good Clinical Practice Guidelines (GCPs) and with the recommendations of the Declaration of Helsinki for investigations with human subjects. The study protocol had been approved by the relevant Ethical Board

(Comitato Etico Regione Calabria, Sezione Area Nord, n° 66/2016) and recorded in a public register of clinical trials (<https://www.clinicaltrials.gov>, NCT03357705).

Every patient signed an informed consent form to document the comprehension of the protocol and of the aims of this study (clinical procedures and potential risks involved). The patient had the possibility to ask questions concerning the treatment and the study protocol and was thoroughly informed about alternative therapies.

A meeting among the clinical centers was held before starting patient recruitment in order to illustrate the protocol and standardize surgical procedures. One trained oral surgeon (with more than 20 years of clinical experience) was selected in each center to perform surgical procedures, whose operative sequence was thoroughly described. The individual responsible for each clinical center received written information to standardize data collection and ensure reliable outcome reporting.

The aims of this parallel-group, multicenter prospective case-control study were the comparison of the dimensional changes of alveolar ridge and sinus floor after maxillary molar extraction with or without performing ARP.

2.2. Study Population. All patients who were 18 years or older and able to sign an informed consent form were considered eligible to participate in this study. Patients underwent a thorough clinical examination to evaluate the state of dentition, including periodontal and occlusal parameters, and a comprehensive treatment plan was discussed and accepted.

2.2.1. Inclusion Criteria

- (i) Indication for the extraction of a first maxillary molar presenting three separated roots;
- (ii) Presence of the adjacent teeth (second premolar and second molar);
- (iii) Presence of intact buccal and palatal bone walls (probing depth ≤ 3 mm);
- (iv) Absence of apical lesions with diameter > 3 mm or cysts.

2.2.2. Exclusion Criteria

- (i) Acute myocardial infarction within the past 2 months;
- (ii) Uncontrolled coagulation disorders;
- (iii) Uncontrolled diabetes (HbA1c $> 7.5\%$);
- (iv) Radiotherapy to the head/neck area within the past 24 months;
- (v) Immunocompromised patients (HIV infection or chemotherapy within the past 5 years);
- (vi) Present or past treatment with intravenous bisphosphonates;
- (vii) Allergy to bovine collagen;

- (viii) Psychological or psychiatric disorders;
- (ix) Alcohol or drugs abuse;
- (x) Full mouth plaque score >30% and/or full mouth bleeding score >20%;
- (xi) Necessity to perform ostectomy procedures to complete dental extraction.

2.3. Treatment Allocation. After a thorough discussion of the various treatment options, patients in whom an implant-supported rehabilitation was planned to replace maxillary first molar were assigned to the test group (ARP). Patients in whom a dental-supported fixed prosthesis was selected to replace the missing molar were assigned to the control group (unassisted socket healing). Patient assignment to the different groups was enclosed in identical, opaque, sealed envelopes which were opened after tooth extraction to reveal to the surgeon the treatment to be performed. Therefore, treatment allocation was concealed to the investigators in charge of enrolling and treating the patients.

2.4. Surgical Procedures. Patients were asked to rinse with chlorhexidine mouthwash 0.2% for 30 seconds. Under local anesthesia (Artin, Omnia, Fidenza, Italy-articain 4% with adrenaline 1:100.000), a mucoperiosteal flap was reflected and minimally traumatic extraction of the tooth was performed after separating roots with ultrasonic tips (Ninja, Acteon, Merignac, France). Roots were then mobilized by micro-elevators and individually extracted by forceps, without performing ostectomy. A careful socket debridement was performed with ultrasonic and manual instruments from the bottom of the socket up to the gingival margin, followed by an accurate removal of the sulcular epithelium. In test group (A), socket was grafted by synthetic nano-hydroxyapatite granules with 250–500 μm diameter (Fisiograft Bone Granular, Ghimas, Casalecchio di Reno, Italy) and covered by haemostatic sponges with collagen of bovine origin (Hemocollagene, Septodont, Saint Maur des Fosses, France); in control group (B) socket was left to spontaneous healing without the insertion of grafting material. In both groups, flaps were mobilized with a periosteal longitudinal releasing incision and sutured with Sentineri technique [27] and single stitches reaching primary closure in both groups. Patients were prescribed with nonsteroidal antiinflammatory drugs (ibuprofen 600 mg), when needed. Sutures were removed after ten days and patients entered in a follow-up protocol with periodic professional dental hygiene recalls. After six months of healing, patients were rehabilitated with dental or implant-supported prostheses according to the previously selected treatment plan.

2.5. Radiographic Examinations. The Ethical Board required the utilization of last generation cone beam computed tomography (CBCT) technology with maximum field of view (FOV) of 5×5 cm, in order to minimize radiation exposure. CBCTs were performed in the area of interest immediately after the end of the surgical procedure (T0) and after six months of healing (T1).




FIGURE 1: Two CBCTs of test group superimposed by Amira software, showing bone present at T0 or at T1 (A1 and B1, respectively); soft tissue present at T0 or at T1 (A2 and B2, respectively), and bone and soft tissues present both at T0 and T1 (C1 and C2, respectively). Schneiderian membrane hypertrophy appears considerably reduced during the healing period.

2.6. Quantitative Radiographic Measurements. T0 and T1 CBCTs of each patient were uploaded on an advanced 3-D image processing and quantification software (Amira, Thermo Fisher Scientific, Waltham, USA). Fiji open-source software [28] was used to perform linear measurements after superimposing CBCTs as previously described by Ryckman et al. [29] (Figure 1). In detail, CBCTs were superimposed by using dental and osseous structures unaffected by the surgery as landmarks (adjacent teeth, lateral and medial maxillary sinus walls, and palatine process): after selecting these regions, the automatic affine registration tool of the software was used for stepwise superimposition. All measurements were taken by a single-blinded calibrated examiner (DP) on a 30-inch led-backlit colour diagnostic display and each measurement was repeated three times at three different time points as proposed by Gomez-Roman and Launer [30]. Examiner calibration was performed by assessing five CBCTs, with another author (FB) who served as reference examiner. Intraexaminer and interexaminer concordances were 95.5% and 91.7%, respectively, for linear measurements within ± 0.1 mm.

In detail, entity of maxillary sinus expansion between T0 and T1 was measured (mm) in correspondence to the apex of the three sockets, following the root axis (PNp, PNd, and PNm) and in the center of the crest (PN). Vertical reduction of the alveolar bone was measured (mm) in correspondence to each of the three sockets, following the root axis, and represented the linear differences between the most coronal position of the ridge at T0 and T1 and the root apex at T0 (RHp, RHd, RHm). Furthermore, alveolar bone height (RH) was measured in the center of the crest (mm), from the most coronal part of the ridge to the sinus floor, both at T0 and at T1. Finally, horizontal reduction of the ridge (RW) was the difference (mm) between bone width in the most coronal part of the crest at T0 (RW0) and T1 (RW1). Figure 2 summarized the reference points taken for measurements.

TABLE 1: Demographic characteristics.

Gender	Age			Sample size		
	Test	Control	Diff.	Test	Control	Diff.
Males	53.4 ± 7.3	51.9 ± 6.5	0.342; NS	7	7	1; NS
Females	56.1 ± 4.2	55.4 ± 5.8		8	8	

Age is presented as mean±standard deviation. Diff., significance of the difference between the groups. NS, no significant difference.




FIGURE 2: Diagram summarizing main reference points taken for measurements. **PN**: sinus expansion between T0 and T1 (mm). **RHm**: vertical reduction of the alveolar bone measured (mm) in correspondence of the mesial socket, following the root axis and representing the linear difference between the most coronal position of the ridge at T0 and T1 and the root apex at T0. **RW0**: ridge width in the most coronal part of the crest at T0. **RW1**: ridge width in the most coronal part of the crest at T1.

2.7. Outcomes. This study evaluated the following outcome measures:

Primary Outcomes

- (i) Maxillary sinus pneumatization (mm): mean of 4 measurements of sinus floor expansion (PNm, PNd, PNp, and PN) from T0 to T1;
- (ii) Alveolar bone dimensional changes (mm): vertical (mean among RHm, RHd, and RHp; RH) and horizontal reduction (RW) of the alveolar bone from T0 to T1.

Secondary Outcomes

- (i) Biological complications: any complication defined as an unexpected deviation from the normal treatment outcome (e.g., alveolitis, postoperative infections).

2.8. Sample Size and Statistical Power. The calculation was performed with a specific software (DSS Research, Fort Worth, USA) to detect a significant difference between the groups in terms of maxillary sinus pneumatization, based on the outcomes of previous research (0.5 mm with an expected standard deviation of 0.3 mm) [26]. A sample of 14 patients (7 test and 7 control cases) was needed to reach 80% of statistical power with α set at 0.05. Each clinical center treated 10 patients for a total of 30 (15 test, 15 control) to

compensate eventual drop-outs occurring during the follow-up period.

2.9. Statistical Analysis. Statistical analysis was performed by means of OriginLab software (OriginLab Corporation, Northampton, USA). Data for descriptive statistics were expressed as mean±SD. All data (with one exception) satisfied both the normality (Kolmogorov–Smirnov test) and the equality of variance (Levene test) assumptions and were analyzed with independent samples T-test. Only RHm of both test and control groups after six months of healing did not satisfy normality and equality of variance and were analyzed by Mann–Whitney test. Fisher exact test was used to evaluate age and gender distribution between the groups. Statistical significance was preset at $\alpha=0.05$.

3. Results

Thirty consecutive patients (14 males and 16 females, age range between 27 and 75, mean 52.7 ± 7.4 , 3 smokers, and 27 nonsmokers) were enrolled and treated with maxillary first molar extraction followed (n=15) or not followed (n=15) by ARP procedures. Complete demographic characteristics of the sample are listed in Table 1: in particular, test and control groups were balanced for age and gender.

Surgeries were performed by three experienced operators [TL (Center 1) n=10; FaB (Center 2) n=10; CS (Center 3) n=10] between November 2016 and June 2017. Four patients (test n=2; control n=2) dropped out at 6-month follow-up (one patient moved abroad, three patients did not come to the control visit within the six month after tooth extraction). Twenty-six patients (test n=13; control n=13) were included in the final analysis. A flowchart diagram summarizing patient selection process was presented in Figure 3.

Six months after dental extraction, sinus floor expansion (mean of PNm, PNd, PNp, and PN) was 0.69 ± 0.48 mm in the test group and 1.04 ± 0.67 mm in the control group ($p=0.15$). Complete results are reported in Table 2.

Vertical resorption of the alveolar bone (mean of RHm, RHd, and RHp) after six months of healing was 1.62 ± 0.49 mm in the test group and 2.01 ± 0.84 mm in the control group ($p=0.08$). Complete data are listed in Table 3.

Available bone height (RH) at T0 was 8.34 ± 3.25 mm and 6.40 ± 1.64 mm in test and control group, respectively ($p=0.07$). At T1, RH was 8.01 ± 3.49 mm and 5.34 ± 2.11 mm in test and control group, respectively ($p=0.03$). Mean vertical bone loss after six months of healing was 0.33 ± 1.94




FIGURE 3: Selection process of patients participating in this study.

mm in test group and 1.06 ± 0.93 mm in control group ($p=0.23$).

Alveolar bone width (RW) at T0 was 11.27 ± 1.71 mm and 12.06 ± 2.46 mm in test and control group, respectively ($p=0.35$). At T1, RW was 8.54 ± 1.26 mm and 8.43 ± 2.26 mm in test and control group, respectively ($p=0.89$). Mean horizontal resorption of crestal bone after six months of healing was 2.73 ± 1.68 mm in test group and 3.63 ± 2.24 mm in control group ($p=0.24$).

No biological complications were recorded during the healing phase. After six months, thirteen patients of the test group underwent dental implant insertion: twelve patients underwent standard implant placement (92.3%); sinus floor elevation was performed in one patient in order to allow implant insertion (7.7%).

4. Discussion

To the best of our knowledge, this paper represents the first prospective case-control study evaluating, with three-dimensional analysis, the clinical effectiveness of ARP in reducing alveolar ridge resorption and maxillary sinus pneumatization after first molar extraction when compared to unassisted socket healing. Previous studies on this topic consisted in retrospective studies based on bidimensional radiographs (orthopantomographies) [23, 25, 26], which do not provide images in bucco-palatal cross section and could be hampered by image distortion [31]. In the present prospective study, measurements were performed by using a semi-automatic method for the comparison of three-dimensional images basing on CBCT superimposition, introduced for

TABLE 2: Entity of sinus pneumatization six months after dental extraction.

	Control (mm)	Test (mm)	Difference (mm)	Diff.
PNm	0.82 ± 0.38	0.59 ± 0.43	0.23	0.15; NS
PNd	0.79 ± 0.71	0.55 ± 0.51	0.24	0.33; NS
PNp	1.08 ± 1.03	0.72 ± 0.70	0.36	0.3; NS
PN	1.46 ± 0.93	0.92 ± 0.63	0.54	0.1; NS
Total	1.04±0.67	0.69±0.48	0.35	0.15; NS

Measures are presented as mean±standard deviation. **PNm**, **PNd**, **PNp**, **PN**, sinus expansion between T0 and T1 measured following the axis of the mesial (**PNm**), distal (**PNd**), palatal (**PNp**) roots and in the center of the crest (**PN**). **Diff.** significance of the difference between the groups. **NS**, no significant difference.

TABLE 3: Vertical resorption of the alveolar bone six months after dental extraction.

	Control (mm)	Test (mm)	Difference (mm)	Diff.
RHm	2.52 ± 1.52	1.45 ± 0.64	1.07	0.20; NS
RHd	1.75 ± 1.24	1.74 ± 0.57	0.01	0.97; NS
RHp	2.31 ± 1.50	1.67 ± 1.02	0.64	0.14; NS
Total	2.01 ± 0.84	1.62 ± 0.49	0.39	0.08; NS

Measures are presented as mean±standard deviation. **RHm**, **RHd**, **RHp**, vertical reduction of the alveolar bone expressed as the difference between the most coronal positions of the ridge at T0 and T1 and the root apex at T0, measured following the axis of the mesial (**RHm**), distal (**RHd**) and palatal (**RHp**) roots. **Diff.** significance of the difference between the groups. **NS**, no significant difference.

the evaluation of bone changes after ARP by Clozza et al. [32]. The accuracy of this procedure is directly dependent on the precision of the superimposition of baseline and final CBCTs: however, possible errors (related to scan acquisition or to incorrect evaluation of the stable regions) are usually negligible, and maxillary regional CBCT superimposition is currently considered as an accurate and reliable method [33].

In the present study, mean extent of sinus pneumatization measured in spontaneously healed sockets six months after tooth extraction was 1.04±0.67 mm, in substantial agreement with previous works by Sharan and Madjar (1.83±2.46 mm) [25] and Levi et al. (1.30±0.27 mm) [26]. Conversely, a smaller downward expansion of the sinus floor, even if not significant, was observed in sites treated with ARP (0.69±0.48 mm), in accordance with the general trend reported by Levi et al. (0.30±0.10 mm) [26].

Hence, these findings suggested that ARP performed after maxillary molar extraction could be an effective technique in reducing maxillary sinus floor pneumatization: mean difference in sinus expansion between test and control group was 0.35 mm after six months of healing. Also this outcome is in accordance with the study of Levi et al. [26], who reported a protective action of ARP on the extent of postextractive sinus pneumatization, even if with a significantly higher mean difference between test and control groups (1 mm).

Postextractive bone remodeling is a widely studied phenomenon regulated by a series of biological events mainly involving bundle bone resorption and resulting in reduced height and width of the residual alveolar ridge [2]. Many clinical trials and systematic reviews with meta-analysis demonstrated that, in general, both horizontal and vertical resorption are more pronounced on the buccal side [34, 35] and that the horizontal reduction (weighted mean 3.87

mm) is greater than the loss in height (weighted mean 1.53 mm) [36]. Alveolar socket grafting was first proposed in the mid-1970s in the attempt to reduce the need of subsequent ridge augmentation procedures prior to implant placement [37, 38]. Many ARP techniques have been proposed lately, improving and extensively testing with various different surgical approaches and biomaterials. Recent systematic reviews with meta-analysis confirmed the effectiveness of ARP in reducing postextractive horizontal and vertical alveolar ridge resorption when compared to spontaneous socket healing, even if the superiority of a specific biomaterial or surgical approach has not been demonstrated yet [17–19, 39–42]. In these studies, the reported clinical magnitude of the effect ranged from 1.4 to 2.19 in terms of buccolingual width and from 1.02 to 2.6 mm in terms of crestal height: the outcomes of the present study (difference between ARP and untreated socket of 0.9 mm in terms of horizontal reduction and 0.39 mm in terms of vertical bone loss) are in line with this trend even if they result in lower values. This occurrence could be possibly explained by the fact that the present study is one of the very few using three-dimensional image processing and measurement tools, which have a higher accuracy and possibility of deeper evaluations than the commonly used bidimensional image analysis systems [43, 44]. Furthermore, the lack of a three-dimensional evaluation could lead to misleading interpretations of particular anatomical conditions, both for clinical and for research purposes (Figure 4).

On this basis, the use of ARP procedures after the extraction of a maxillary molar could be regarded as a preventive treatment, particularly in cases where an implant-supported restoration is planned. Maxillary molars (particularly first molar) are the sites in which the combined action of alveolar bone remodeling and maxillary sinus




FIGURE 4: An example of a particular anatomical situation in which the lack of three-dimensional evaluation could lead to misleading interpretations of available bone height.

pneumatization leads to the greater amount of vertical bone loss after tooth extraction of the entire upper arch [4, 23]. Hence, a preservation of vertical and horizontal alveolar ridge dimensions could often be clinically significant, allowing a standard implant placement without additional regenerative procedures. Sinus floor elevation (both lateral and transcrestal) and GBR techniques are advanced procedures requiring specific operator skills and are always associated with higher morbidity, increasing costs, prolonged time of therapy, and potential intra- and postoperative complications [12, 45–48]. In our test group, mean available crestal height (RH) was 8.34 mm at baseline and 8.01 mm six months after molar extraction, presenting only a 4% reduction: 12 patients out of 13 underwent implant insertion without the need of additional bone augmentation procedures. This finding is in accordance with the results of a randomized trial by Rasperini et al. [49], confirming that ARP performed in the posterior maxilla increases the possibility of inserting implants without the need for contextual sinus augmentation procedures.

Some limitations should be considered when interpreting the outcomes of this study. The first to consider is the strict local inclusion criteria (three separated roots; presence of the adjacent teeth; presence of intact buccal and palatal bone walls): ARP effectiveness should be tested also in different conditions. Secondly, the use of CBCT as three-dimensional data source, even if more reliable than bidimensional radiographs, may be subjected to lack of standardization among different equipment. Third, no histologic evaluations have been performed in the present study: even if the use of synthetic grafts in ARP is supported by recent randomized controlled trials with human histologies [50, 51], further research is necessary to confirm the effectiveness of this class of biomaterials in this particular application.

Finally, randomized clinical trials on larger samples are recommended, in order to prevent potential bias in patient

allocation to the different treatment groups and to reduce data dispersion.

5. Conclusion

Within the limitations of this study, it could be suggested that ARP performed after maxillary molar extraction may reduce the entity of sinus pneumatization and alveolar bone resorption, compared to unassisted socket healing. The possibility of an effective preservation of vertical and horizontal crestal dimensions could decrease the necessity of advanced regenerative procedures prior to dental implant placement: further studies are necessary to confirm these findings and to better define the clinical magnitude of the effect.

Data Availability

The datasets generated and analyzed during the current study are available from the corresponding author upon reasonable request.

Conflicts of Interest

The authors declare no conflicts of interest related to this study.

Acknowledgments

The authors wish to thank Dr. Paolo Ceribelli and Mrs. Romana Villa for their precious help in the conduction of this study and in manuscript revision.

References

- [1] L. Schropp, A. Wenzel, L. Kostopoulos, and T. Karring, "Bone healing and soft tissue contour changes following single-tooth extraction: a clinical and radiographic 12-month prospective study," *International Journal of Periodontics and Restorative Dentistry*, vol. 23, no. 4, pp. 313–323, 2003.
- [2] L. Trombelli, R. Farina, A. Marzola, L. Bozzi, B. Liljenberg, and J. Lindhe, "Modeling and remodeling of human extraction sockets," *Journal of Clinical Periodontology*, vol. 35, no. 7, pp. 630–639, 2008.
- [3] B. K. Bartee, "Extraction site reconstruction for alveolar ridge preservation. Part 2: membrane-assisted surgical technique," *Journal of Oral Implantology*, vol. 27, no. 4, pp. 194–197, 2001.
- [4] M. Pramstraller, R. Farina, G. Franceschetti, C. Pramstraller, and L. Trombelli, "Ridge dimensions of the edentulous posterior maxilla: A retrospective analysis of a cohort of 127 patients using computerized tomography data," *Clinical Oral Implants Research*, vol. 22, no. 1, pp. 54–61, 2011.
- [5] G. Avila, H.-L. Wang, P. Galindo-Moreno et al., "The influence of the bucco-palatal distance on sinus augmentation outcomes," *Journal of Periodontology*, vol. 81, no. 7, pp. 1041–1050, 2010.
- [6] S. Corbella, S. Taschieri, R. Weinstein, and M. Del Fabbro, "Histomorphometric outcomes after lateral sinus floor elevation procedure: a systematic review of the literature and meta-analysis," *Clinical Oral Implants Research*, vol. 27, no. 9, pp. 1106–1122, 2016.

- [7] C. Stacchi, T. Lombardi, R. Ottonelli, F. Berton, G. Perinetti, and T. Traini, "New bone formation after transcrestal sinus floor elevation was influenced by sinus cavity dimensions: A prospective histologic and histomorphometric study," *Clinical Oral Implants Research*, vol. 29, no. 5, pp. 465–479, 2018.
- [8] M. Beretta, M. Cicciù, and P. P. Poli, "A retrospective evaluation of 192 implants placed in augmented bone: long-term follow-up study," *Journal of Oral Implantology*, vol. 41, no. 6, pp. 669–674, 2015.
- [9] G. Orsini, C. Stacchi, E. Visintini et al., "Clinical and histologic evaluation of fresh frozen human bone grafts for horizontal reconstruction of maxillary alveolar ridges," *The International Journal of Periodontics & Restorative Dentistry*, vol. 31, no. 5, pp. 535–544, 2011.
- [10] A. Monje, M. A. Pikos, H.-L. Chan et al., "On the feasibility of utilizing allogeneic bone blocks for atrophic maxillary augmentation," *BioMed Research International*, vol. 2014, Article ID 814578, 12 pages, 2014.
- [11] G. N. Güncü, Y. D. Yildirim, H.-L. Wang, and T. F. Tözüm, "Location of posterior superior alveolar artery and evaluation of maxillary sinus anatomy with computerized tomography: a clinical study," *Clinical Oral Implants Research*, vol. 22, no. 10, pp. 1164–1167, 2011.
- [12] C. Stacchi, F. Andolsek, F. Berton, G. Perinetti, C. O. Navarra, and R. Di Lenarda, "Intraoperative complications during sinus floor elevation with lateral approach: A systematic review," *The International Journal of Oral & Maxillofacial Implants*, vol. 32, no. 3, pp. e107–e118, 2017.
- [13] M. Esposito, M. G. Grusovin, and J. Rees, "Interventions for replacing missing teeth: augmentation procedures of the maxillary sinus," *Cochrane Database of Systematic Reviews*, vol. 17, no. 5, Article ID CD008397, 2014.
- [14] S. Spinato, A. Rebaudi, F. Bernardello, C. Bertoldi, and D. Zaffe, "Piezosurgical treatment of crestal bone: Quantitative comparison of post-extractive socket outcomes with those of traditional treatment," *Clinical Oral Implants Research*, vol. 27, no. 3, pp. 361–366, 2016.
- [15] E. Bramanti, A. Norcia, M. Cicciù et al., "Postextraction dental implant in the aesthetic zone, socket shield technique versus conventional protocol," *The Journal of Craniofacial Surgery*, vol. 29, no. 4, pp. 1037–1041, 2018.
- [16] S. Spinato, P. Galindo-Moreno, D. Zaffe, F. Bernardello, and C. M. Soardi, "Is socket healing conditioned by buccal plate thickness? A clinical and histologic study 4 months after mineralized human bone allografting," *Clinical Oral Implants Research*, vol. 25, no. 2, pp. e120–e126, 2014.
- [17] L. Barallat, V. Ruiz-Magaz, P. A. Levi, S. Mareque-Bueno, P. Galindo-Moreno, and J. Nart, "Histomorphometric results in ridge preservation procedures comparing various graft materials in extraction sockets with nongrafted sockets in humans: A systematic review," *Implant Dentistry*, vol. 23, no. 5, pp. 539–554, 2014.
- [18] G. Troiano, K. Zhurakivska, L. Lo Muzio, L. Laino, M. Cicciù, and L. Lo Russo, "Combination of bone graft and resorbable membrane for alveolar ridge preservation: a systematic review, meta-analysis and trial sequential analysis," *Journal of Periodontology*, vol. 89, no. 1, pp. 46–57, 2018.
- [19] N. Macbeth, A. Trullenque-Eriksson, N. Donos, and N. Mardas, "Hard and soft tissue changes following alveolar ridge preservation: a systematic review," *Clinical Oral Implants Research*, vol. 28, no. 8, pp. 982–1004, 2017.
- [20] B. Leblebicioglu, M. Salas, Y. Ort et al., "Determinants of alveolar ridge preservation differ by anatomic location," *Journal of Clinical Periodontology*, vol. 40, no. 4, pp. 387–395, 2013.
- [21] D. Cardaropoli, L. Tamagnone, A. Roffredo, L. Gaveglio, and G. Cardaropoli, "Socket preservation using bovine bone mineral and collagen membrane: a randomized controlled clinical trial with histologic analysis," *The International Journal of Periodontics & Restorative Dentistry*, vol. 32, no. 4, pp. 421–430, 2012.
- [22] A. Barone, M. Ricci, P. Tonelli, S. Santini, and U. Covani, "Tissue changes of extraction sockets in humans: a comparison of spontaneous healing vs. ridge preservation with secondary soft tissue healing," *Clinical Oral Implants Research*, vol. 24, no. 11, pp. 1231–1237, 2013.
- [23] H. Wehrbein and P. Diedrich, "Progressive pneumatization of the basal maxillary sinus after extraction and space closure," *Fortschritte der Kieferorthopädie*, vol. 53, no. 2, pp. 77–83, 1992.
- [24] H. M. Frost, "Wolff's Law and bone's structural adaptations to mechanical usage: an overview for clinicians," *The Angle Orthodontist*, vol. 64, no. 3, pp. 175–188, 1994.
- [25] A. Sharan and D. Madjar, "Maxillary sinus pneumatization following extractions: a radiographic study," *The International Journal of Oral & Maxillofacial Implants*, vol. 23, no. 1, pp. 48–56, 2008.
- [26] I. Levi, M. Halperin-Sternfeld, J. Horwitz, H. Zigdon-Giladi, and E. E. Machtei, "Dimensional changes of the maxillary sinus following tooth extraction in the posterior maxilla with and without socket preservation," *Clinical Implant Dentistry and Related Research*, vol. 19, no. 5, pp. 952–958, 2017.
- [27] R. Sentineri, T. Lombardi, F. Berton, and C. Stacchi, "Laurell-Gottlow suture modified by Sentineri for tight closure of a wound with a single line of sutures," *British Journal of Oral and Maxillofacial Surgery*, vol. 54, no. 1, pp. e18–e19, 2016.
- [28] J. Schindelin, I. Arganda-Carreras, and E. Frise, "Fiji: an open-source platform for biological-image analysis," *Nature Methods*, vol. 9, no. 7, pp. 676–682, 2012.
- [29] M. S. Ryckman, S. Harrison, D. Oliver et al., "Soft-tissue changes after maxillomandibular advancement surgery assessed with cone-beam computed tomography," *American Journal of Orthodontics and Dentofacial Orthopedics*, vol. 137, no. 4, pp. S86–S93, 2010.
- [30] G. Gomez-Roman and S. Launer, "Peri-implant bone changes in immediate and non-immediate root-analog stepped implants—a matched comparative prospective study up to 10 years," *International Journal of Implant Dentistry*, vol. 2, no. 1, p. 15, 2016.
- [31] F. Allen and D. G. Smith, "An assessment of the accuracy of ridge-mapping in planning implant therapy for the anterior maxilla," *Clinical Oral Implants Research*, vol. 11, no. 1, pp. 34–38, 2000.
- [32] E. Clozza, M. Biasotto, F. Cavalli, L. Moimas, and R. Di Lenarda, "Three-dimensional evaluation of bone changes following ridge preservation procedures," *The International Journal of Oral & Maxillofacial Implants*, vol. 27, no. 4, pp. 770–775, 2012.
- [33] L. Koerich, D. Burns, A. Weissheimer, and J. D. P. Claus, "Three-dimensional maxillary and mandibular regional superimposition using cone beam computed tomography: a validation study," *International Journal of Oral and Maxillofacial Surgery*, vol. 45, no. 5, pp. 662–669, 2016.
- [34] V. Chappuis, O. Engel, M. Reyes, K. Shahim, L.-P. Nolte, and D. Buser, "Ridge alterations post-extraction in the esthetic zone: a

- 3D analysis with CBCT,” *Journal of Dental Research*, vol. 92, no. 12, pp. 1955–2015, 2013.
- [35] M. Misawa, J. Lindhe, and M. G. Araújo, “The alveolar process following single-tooth extraction: a study of maxillary incisor and premolar sites in man,” *Clinical Oral Implants Research*, vol. 27, no. 7, pp. 884–889, 2016.
- [36] F. Van der Weijden, F. Dell’Acqua, and D. E. Slot, “Alveolar bone dimensional changes of post-extraction sockets in humans: a systematic review,” *Journal of Clinical Periodontology*, vol. 36, no. 12, pp. 1048–1058, 2009.
- [37] A. C. De Carvalho, M. C. Pinto, and V. de Sousa, “Implant of “synthetic bone” in the dental alveolus. Clinical and radiographic study of patients,” *Revista da Associação Paulista de Cirurgiões Dentistas*, vol. 26, no. 3, pp. 125–128, 1972.
- [38] R. Marques-Miguel, M. G. Sanches, and M. C. Santos-Pinto, “Healing of extraction wounds after anorganic bone implant. Radiographic study in rats,” *Revista da Associação Paulista de Cirurgiões Dentistas*, vol. 27, no. 2, pp. 88–93, 1973.
- [39] K. R. Morjaria, R. Wilson, and R. M. Palmer, “Bone healing after tooth extraction with or without an intervention: A systematic review of randomized controlled trials,” *Clinical Implant Dentistry and Related Research*, vol. 16, no. 1, pp. 1–20, 2014.
- [40] G. Avila-Ortiz, S. Elangovan, K. W. O. Kramer, D. Blanchette, and D. V. Dawson, “Effect of alveolar ridge preservation after tooth extraction: a systematic review and meta-analysis,” *Journal of Dental Research*, vol. 93, no. 10, pp. 950–958, 2014.
- [41] M. A. Atieh, N. H. Alsabeeha, A. G. Payne, W. Duncan, C. M. Faggion, and M. Esposito, “Interventions for replacing missing teeth: Alveolar ridge preservation techniques for dental implant site development,” *Cochrane Database of Systematic Reviews*, vol. 2015, no. 5, 2015.
- [42] O. Iocca, A. Farcomeni, S. Pardiñas Lopez, and H. S. Talib, “Alveolar ridge preservation after tooth extraction: a Bayesian Network meta-analysis of grafting materials efficacy on prevention of bone height and width reduction,” *Journal of Clinical Periodontology*, vol. 44, no. 1, pp. 104–114, 2017.
- [43] W. M. Takeshita, L. C. V. Iwaki, M. C. Da Silva, and R. H. Tonin, “Evaluation of diagnostic accuracy of conventional and digital periapical radiography, panoramic radiography, and cone-beam computed tomography in the assessment of alveolar bone loss,” *Contemporary Clinical Dentistry*, vol. 5, no. 3, pp. 318–323, 2014.
- [44] P. Pittayapat, M. M. Bornstein, T. S. N. Imada, W. Coucke, I. Lambrichts, and R. Jacobs, “Accuracy of linear measurements using three imaging modalities: Two lateral cephalograms and one 3D model from CBCT data,” *European Journal of Orthodontics*, vol. 37, no. 2, pp. 202–208, 2014.
- [45] F. Fontana, E. Maschera, I. Rocchietta, and M. Simion, “Clinical classification of complications in guided bone regeneration procedures by means of a nonresorbable membrane,” *International Journal of Periodontics and Restorative Dentistry*, vol. 31, no. 3, pp. 265–273, 2011.
- [46] P. Fugazzotto, P. R. Melnick, and M. Al-Sabbagh, “Complications when augmenting the posterior maxilla,” *Dental Clinics of North America*, vol. 59, no. 1, pp. 97–130, 2015.
- [47] C. Stacchi, R. Sentineri, F. Berton, and T. Lombardi, “Conjunctival chemosis: an uncommon complication after transcrestal lifting of the sinus floor,” *British Journal of Oral and Maxillofacial Surgery*, vol. 54, no. 9, pp. 1052–1054, 2016.
- [48] R. Farina, A. Zaetta, L. Minenna, and L. Trombelli, “Orbital and periorbital emphysema following maxillary sinus floor elevation: a case report and literature review,” *Journal of Oral and Maxillofacial Surgery*, vol. 74, no. 11, pp. 2192–2192.e7, 2016.
- [49] G. Rasperini, L. Canullo, C. Dellavia, G. Pellegrini, and M. Simion, “Socket grafting in the posterior maxilla reduces the need for sinus augmentation,” *International Journal of Periodontics and Restorative Dentistry*, vol. 30, no. 3, pp. 265–273, 2010.
- [50] G. A. Gholami, B. Najafi, F. Mashhadiabbas, W. Goetz, and S. Najafi, “Clinical, histologic and histomorphometric evaluation of socket preservation using a synthetic nanocrystalline hydroxyapatite in comparison with a bovine xenograft: A randomized clinical trial,” *Clinical Oral Implants Research*, vol. 23, no. 10, pp. 1198–1204, 2012.
- [51] Y. Mayer, H. Zigdon-Giladi, and E. E. Machtei, “Ridge preservation using composite alloplastic materials: a randomized control clinical and histological study in humans,” *Clinical Implant Dentistry and Related Research*, vol. 18, no. 6, pp. 1163–1170, 2016.